



Stability Analysis for the Gurtin-MacCamy's Age-Structured Population Dynamics Model

Mohammed El-doma

Faculty of Mathematical Sciences
University of Khartoum
P. O. Box 321, Khartoum, Sudan
E-mail: biomath2004@yahoo.com
Telephone: 249183774443, Fax: 249183780295

Received February 12,2007; accepted November 7, 2007

Abstract

The stability of the Gurtin-MacCamy's age-structured population dynamics model is investigated. We determine the steady states and study their stability. The results in this paper generalize previous results.

Keywords: Population; Stability; Steady state; Age-structure.

MSC 2000: 45K05; 45M05; 45M10; 45M20; 35B35; 35B40; 35L60; 92D25.

1. Introduction

In this paper, we study the Gurtin-MacCamy's age-structured population dynamics model that has been studied in Gurtin, et al. (1974). The model is as follows

$$\begin{cases} \frac{\partial p(a,t)}{\partial a} + \frac{\partial p(a,t)}{\partial t} + \mu(a, P(t))p(a,t) = 0, & a > 0, \quad t > 0, \\ p(0,t) = \int_0^{\infty} \beta(a, P(t))p(a,t)da, & t \geq 0, \\ p(a,0) = p_0(a), & a \geq 0, \\ P(t) = \int_0^{\infty} p(a,t)da, & t \geq 0, \end{cases} \quad (1)$$

where $p(a,t)$ is the density of the population with respect to the chronological age $a \in [0, \infty)$

at time $t \geq 0$; $P(t) = \int_0^\infty p(a, t) da$ is the total population size at time t ; $\beta(a, P(t)), \mu(a, P(t))$ are, respectively, the birth rate i.e. the average number of offspring, per unit time, produced by an individual of age a when the population size is $P(t)$, and the mortality rate i.e. the death rate at age a , per unit population, when the population size is $P(t)$; $p(0, t) = \int_0^\infty \beta(a, P(t))p(a, t) da$ is the number of births, per unit time, when the population size is $P(t)$.

We study problem (1) under the following general assumptions:

$$\left\{ \begin{array}{l} 0 \leq p_0(a) \in L^1(\mathbb{R}^+), \quad \mathbb{R}^+ = [0, \infty); \\ \beta(a, P(t)) > 0, \quad \mu(a, P(t)) > 0, \quad \mu(a, P(t)) \in C(\mathbb{R}^+ \times \mathbb{R}^+); \\ \beta_P(a, P), \quad \mu_P(a, P) \text{ exist for } \forall a \geq 0, P \geq 0; \\ \beta(\cdot, P), \beta_P(\cdot, P), \mu(\cdot, P), \mu_P(\cdot, P) \in C(\mathbb{R}^+; L_\infty(\mathbb{R}^+)). \end{array} \right. \quad (2)$$

In Gurtin, et al. (1974) problem (1) is described and existence and uniqueness results via a fixed point theorem are obtained as well as the local asymptotic stability of a nontrivial steady state when all the roots of the corresponding characteristic equation have negative real parts. Since then several authors have attempted to study the stability of special cases of this model in order to determine conditions under which all the roots of the corresponding characteristic equation have negative real parts, for example, see Weinstock, et al. (1987), Di Blasio, et al. (1982), and Iannelli(1995) and the references therein. Some recent results involving the global stability and global boundedness of solution are given in Iannelli(1995), Iannelli, et al. (2002), Bekkal-Brikci, et al. (2007), and El-Doma (Preprint to appear).

In this paper, we study problem (1) and determine its steady states and examine their stability. We show that if $R(0) < 1$, see section 2 for the definition of $R(P)$, then the trivial steady state is locally asymptotically stable, and if $R(0) > 1$, then the trivial steady state is unstable. We also determine sufficient conditions for the local asymptotic stability of a nontrivial steady state P_∞ , and show that if $R'(P_\infty) > 0$, then a nontrivial steady state is unstable.

The organization of this paper as follows: in section 2 we determine the steady states; in section 3 we study the stability of the steady states; in section 4 we conclude our results.

2. The Steady States

In this section, we determine the steady states of problem (1). A steady state of problem (1) satisfies the following:

$$\left\{ \begin{array}{l} \frac{dp_\infty(a)}{da} + \mu(a, P_\infty)p_\infty(a) = 0, \quad a > 0, \\ p_\infty(0) = \int_0^\infty \beta(a, P_\infty)p_\infty(a) da, \\ P_\infty = \int_0^\infty p_\infty(a) da. \end{array} \right. \quad (3)$$

From (3), by solving the differential equation, we obtain that

$$p_\infty(a) = p_\infty(0)\pi(a, P_\infty), \quad (4)$$

where $\pi(a, P_\infty)$ is defined as

$$\pi(a, P) = e^{-\int_0^a \mu(\tau, P) d\tau},$$

is the probability of survival to age a when the population size is P . Also, from (3) and (4), we obtain that $p_\infty(0)$ satisfies the following:

$$p_\infty(0) = p_\infty(0) \int_0^\infty \beta(a, P_\infty) \pi(a, P_\infty) da. \quad (5)$$

Accordingly, from (5), we conclude that either $p_\infty(0) = 0$ or P_∞ satisfies the following:

$$1 = \int_0^\infty \beta(a, P_\infty) \pi(a, P_\infty) da. \quad (6)$$

In order to facilitate our writing, we define a threshold parameter $R(P)$ by

$$R(P) = \int_0^\infty \beta(a, P) \pi(a, P) da, \quad (7)$$

which is interpreted as the number of children expected to be born to an individual, in a life time, when the population size is P .

We note that from equation (4), $p_\infty(0) = \frac{P_\infty}{\int_0^\infty \pi(a, P_\infty) da}$, and accordingly, either $p_\infty(a) \equiv 0$

or $p_\infty(a)$ is completely determined by a solution P_∞ of equation (6).

In the following theorem, we describe the steady states of problem (1), the proof of the theorem is straightforward and therefore, is omitted.

Theorem 2.1:

- (a) Problem (1) has the trivial steady state, $P_\infty = 0$, as a steady state.
- (b) All nontrivial solutions of, $R(P_\infty) = 1$, are nontrivial steady states of problem (1).

3. Stability of the Steady States

In this section, we study the stability of the steady states for problem (1) as given by Theorem 2.1.

To study the stability of a steady state $p_\infty(a)$, which is a solution of (3) and is given by equation (4), we linearize problem (1) at $p_\infty(a)$ in order to obtain a characteristic equation, which in turn will determine conditions for the stability. To that end, we consider a perturbation $\omega(a, t)$ defined by $\omega(a, t) = p(a, t) - p_\infty(a)$, where $p(a, t)$ is a solution of problem (1). Accordingly, we obtain that $\omega(a, t)$ satisfies the following:

$$\begin{cases} \frac{\partial \omega(a, t)}{\partial a} + \frac{\partial \omega(a, t)}{\partial t} + \mu(a, P_\infty) \omega(a, t) + p_\infty(a) \mu_P(a, P_\infty) W(t) = 0, & a > 0, \quad t > 0, \\ \omega(0, t) = \int_0^\infty \beta(a, P_\infty) \omega(a, t) da + W(t) \int_0^\infty \beta_P(a, P_\infty) p_\infty(a) da, & t \geq 0, \\ \omega(a, 0) = p_0(a) - p_\infty(a), & a \geq 0, \\ W(t) = \int_0^\infty \omega(a, t) da, & t \geq 0. \end{cases} \quad (8)$$

By substituting $\omega(a, t) = f(a)e^{\lambda t}$ in (8), where λ is a complex number, and straightforward calculations, we obtain the following characteristic equation:

$$1 = \int_0^\infty e^{-\lambda a} \beta(a, P_\infty) \pi(a, P_\infty) da + \frac{\int_0^\infty e^{-\lambda a} \pi(a, P_\infty) da}{1 + \int_0^\infty \int_0^a e^{-\lambda(a-\sigma)} p_\infty(a) \mu_P(\sigma, P_\infty) d\sigma da} \times \left[\int_0^\infty \beta_P(a, P_\infty) p_\infty(a) da - \int_0^\infty \int_0^a e^{-\lambda(a-\sigma)} \beta(a, P_\infty) p_\infty(a) \mu_P(\sigma, P_\infty) d\sigma da \right]. \tag{9}$$

In the following theorem, we describe the stability of the trivial steady state $p_\infty(a) \equiv 0$.

Theorem 3.1: The trivial steady state $p_\infty(a) \equiv 0$, is locally asymptotically stable if $R(0) < 1$, and is unstable if $R(0) > 1$.

Proof: We note that for the trivial steady state $p_\infty(a) \equiv 0$, $P_\infty = 0$, and therefore, from the characteristic equation (9), we obtain the following characteristic equation:

$$1 = \int_0^\infty e^{-\lambda a} \beta(a, 0) \pi(a, 0) da. \tag{10}$$

To prove the local asymptotic stability of the trivial steady state, we note that if $R(0) < 1$, then equation (10) can not be satisfied for any λ with $Re\lambda \geq 0$ since

$$\left| \int_0^\infty \beta(a, 0) \pi(a, 0) e^{-\lambda a} da \right| \leq \int_0^\infty \beta(a, 0) \pi(a, 0) e^{-a Re\lambda} da, \leq R(0) < 1.$$

Accordingly, the trivial steady state is locally asymptotically stable if $R(0) < 1$.

To prove the instability of the trivial steady state when $R(0) > 1$, we note that if we define a function $g(\lambda)$ by

$$g(\lambda) = \int_0^\infty \beta(a, 0) \pi(a, 0) e^{-\lambda a} da,$$

and suppose that λ is real, then we can easily see that $g(\lambda)$ is a decreasing function if $\lambda > 0$, $g(\lambda) \rightarrow 0$ as $\lambda \rightarrow +\infty$, and $g(0) = R(0)$. Therefore, if $R(0) > 1$, then there exists $\lambda^* > 0$ such that $g(\lambda^*) = 1$, and hence the trivial steady state is unstable. This completes the proof of the theorem.

In the next theorem, we give a condition for the instability of a nontrivial steady state.

Theorem 3.2: A nontrivial steady state is unstable if $R'(P_\infty) > 0$.

Proof: We note that the characteristic equation (9) can be rewritten as

$$\begin{aligned}
1 &= \int_0^\infty e^{-\lambda a} \beta(a, P_\infty) \pi(a, P_\infty) da \\
&+ \int_0^\infty \int_0^a e^{-\lambda(a-\sigma)} p_\infty(a) \mu_P(\sigma, P_\infty) d\sigma da \left[\int_0^\infty e^{-\lambda a} \beta(a, P_\infty) \pi(a, P_\infty) da - 1 \right] \\
&+ \int_0^\infty e^{-\lambda a} \pi(a, P_\infty) da \times \\
&\quad \left(\int_0^\infty \beta_P(a, P_\infty) p_\infty(a) da - \int_0^\infty \int_0^a e^{-\lambda(a-\sigma)} \beta(a, P_\infty) p_\infty(a) \mu_P(\sigma, P_\infty) d\sigma da \right) \\
&:= G(\lambda).
\end{aligned} \tag{11}$$

Now, suppose that $R'(P_\infty) > 0$, then from the characteristic equation (11), we obtain that $G(0) = 1 + R'(P_\infty)P_\infty > 1$, and $G(\lambda) \rightarrow 0$ as $\lambda \rightarrow +\infty$. Accordingly, $\exists \lambda^* > 0$ such that $G(\lambda^*) = 1$, and hence a nontrivial steady state is unstable. This completes the proof of the theorem.

In the next theorem, we prove that, $\lambda = 0$, is a root of the characteristic equation (11) iff $R'(P_\infty) = 0$.

Theorem 3.3: $\lambda = 0$ is a root of the characteristic equation (11) iff $R'(P_\infty) = 0$.

Proof: We note that if $\lambda = 0$, then using equation (6), the characteristic equation (11) becomes $R'(P_\infty) = 0$. This completes the proof of the theorem.

To obtain further stability results, we note that by suitable changes of the variables of the integrations, we can rewrite the characteristic equation (11) in the following form:

$$\begin{aligned}
1 &= \int_0^\infty e^{-\lambda a} \left\{ \beta(a, P_\infty) \pi(a, P_\infty) - \int_a^\infty p_\infty(\sigma) \mu_P(\sigma - a, P_\infty) d\sigma \right. \\
&+ \pi(a, P_\infty) \int_0^\infty \beta_P(b, P_\infty) p_\infty(b) db \\
&\left. + \int_0^a \int_{a-b}^\infty p_\infty(b) \pi(\sigma, P_\infty) \mu_P(\sigma + b - a, P_\infty) [\beta(b, P_\infty) - \beta(\sigma, P_\infty)] d\sigma db \right\} da.
\end{aligned} \tag{12}$$

In the next theorem, we give our first sufficient condition for the local asymptotic stability of a nontrivial steady state. We note that contrary to our results in this paper, most stability results are obtained for special cases of problem (1), for example, see Weinstock, et al. (1987), Di Blasio, et al. (1982), and Iannelli(1995) and the references therein.

Theorem 3.4: Suppose that the following hold:

$$\begin{aligned}
1. \quad \phi(a, P_\infty) &= \beta(a, P_\infty) \pi(a, P_\infty) - \int_a^\infty p_\infty(\sigma) \mu_P(\sigma - a, P_\infty) d\sigma \\
&+ \pi(a, P_\infty) \int_0^\infty \beta_P(b, P_\infty) p_\infty(b) db \\
&+ \int_0^a \int_{a-b}^\infty p_\infty(b) \pi(\sigma, P_\infty) \mu_P(\sigma + b - a, P_\infty) [\beta(b, P_\infty) - \beta(\sigma, P_\infty)] d\sigma db \geq 0, \\
&\forall a \geq 0,
\end{aligned} \tag{13}$$

2. $R'(P_\infty) < 0$.

Then a nontrivial steady state is locally asymptotically stable.

Proof: We note that from the characteristic equation (12), we obtain that

$$1 = \int_0^\infty e^{-\lambda a} \phi(a, P_\infty) da. \tag{14}$$

And therefore, if we assume that $Re\lambda \geq 0$, then from equation (14), we obtain

$$\begin{aligned} \left| \int_0^\infty e^{-\lambda a} \phi(a, P_\infty) da \right| &\leq \int_0^\infty e^{-aRe\lambda} \phi(a, P_\infty) da \\ &\leq \int_0^\infty \phi(a, P_\infty) da \\ &= 1 + R'(P_\infty)P_\infty \\ &< 1. \end{aligned}$$

Accordingly, the characteristic equation (12) is not satisfied for any λ with real $Re\lambda \geq 0$, and hence a nontrivial steady state is locally asymptotically stable. This completes the proof of the theorem.

In the next result, we give a corollary to Theorem 3.4, and the proof of this corollary is straightforward, and therefore, is omitted.

Corollary 3.5: Suppose that the following hold:

- 1) $\beta(a, P) = \beta(P), \forall a \geq 0$,
- 2) $\pi(a, P_\infty) \left[\beta(P_\infty) + \beta'(P_\infty)P_\infty \right] - \int_a^\infty p_\infty(\sigma) \mu_P(\sigma - a, P_\infty) d\sigma \geq 0, \forall a \geq 0$,
- 3) $R'(P_\infty) < 0$.

Then a nontrivial steady state is locally asymptotically stable.

In the next theorem, we give our second sufficient condition for the local asymptotic stability of a nontrivial steady state.

Theorem 3.6: Suppose that $\int_0^\infty |\phi(a, P_\infty)| da < 1$, where ϕ is given by equation (13). Then a nontrivial steady state is locally asymptotically stable.

Proof: Suppose that $Re\lambda \geq 0$, then from the characteristic equation (12), we obtain

$$\begin{aligned} 1 = \left| \int_0^\infty e^{-\lambda a} \phi(a, P_\infty) da \right| &\leq \int_0^\infty e^{-aRe\lambda} |\phi(a, P_\infty)| da \\ &\leq \int_0^\infty |\phi(a, P_\infty)| da \\ &< 1. \end{aligned}$$

Accordingly, the characteristic equation (12) is not satisfied for any λ with real $Re\lambda \geq 0$, and hence a nontrivial steady state is locally asymptotically stable. This completes the proof of the theorem.

We note that a similar corollary to corollary 3.5 holds in the case when β is a function of P only.

Example: If we consider the example given in Gurtin, et al. (1974), where $\mu(a, P) = \mu(P)$; $\beta(a, P) = \beta(P)e^{-\alpha a}$, $\alpha \geq 0$. Then $R(P) = \frac{\beta(P)}{\mu(P) + \alpha}$. And therefore, $R'(P_\infty) < 0$, is equivalent to $\beta'(P_\infty) < \mu'(P_\infty)$. And in this case Theorem 3.4 gives the following conditions for the local asymptotic stability:

- 1) $\beta(P_\infty)e^{-\beta(P_\infty)a} + p_\infty(0)e^{-\mu(P_\infty)a} \left[\frac{\beta'(P_\infty)}{\beta(P_\infty)} - \frac{\mu'(P_\infty)}{\mu(P_\infty)}e^{-\alpha a} \right] \geq 0$,
- 2) $\beta'(P_\infty) < \mu'(P_\infty)$.

Also by Theorem 3.6 and if we assume the positivity condition:

$$\beta(P_\infty)e^{-\beta(P_\infty)a} + p_\infty(0)e^{-\mu(P_\infty)a} \left[\frac{\beta'(P_\infty)}{\beta(P_\infty)} - \frac{\mu'(P_\infty)}{\mu(P_\infty)}e^{-\alpha a} \right] \geq 0,$$

then we obtain the following condition for the local asymptotic stability:

$$1 + \frac{p_\infty(0)}{\beta(P_\infty)\mu(P_\infty)} [\beta'(P_\infty) - \mu'(P_\infty)] < 1,$$

which clearly satisfies the condition of the theorem when $\beta'(P_\infty) - \mu'(P_\infty) < 0$, which is exactly the condition $R'(P_\infty) < 0$.

4. Conclusion

In this paper, we studied the stability of the well-known Gurtin-MacCamy's age-structured population dynamics model given in their paper Gurtin, et al.(1974). In our results, we obtained stability results for their general model, however, the stability of special cases of this model has been given in several papers and books, for example, see Weinstock, et al.(1987), Di Blasio, et al. (1982), and Iannelli(1995), and the references therein. Some recent results involving the global stability as well as the global boundedness of solution are given in Iannelli(1995), Iannelli, et al. (2002), Bekkal-Brikci, et al.(2007), and El-Doma (Preprint to appear).

We determined the steady states of the model and examined their stability. We proved that the trivial steady state is always a steady state and that there are as many nontrivial steady states as the solutions of the equation, $R(P_\infty) = 1$, where $R(P)$ is the number of children expected to be born to an individual, over a life time, when the population size is P . Then we studied the stability of the trivial steady state and showed that if $R(0) < 1$, then the trivial steady state is locally asymptotically stable, and if $R(0) > 1$, then the trivial steady state is unstable.

Furthermore, we studied the stability of a nontrivial steady state and we gave several sufficient conditions for their local asymptotic stability. We also showed that if $R'(P_\infty) > 0$, then a nontrivial steady state is unstable.

References

Bekkal-Brikci, F., K. Boushaba and O. Arino. Nonlinear age structured model with cannibalism. *Discrete and Continuous Dynamical Systems-Series B*. Vol. 7, pp. 201-218, (2007).

Di Blasio, G., M. Iannelli and E. Sinestrari. Approach to Equilibrium in Age Structured Populations with an Increasing Recruitment Process. *J. Math. Biol.* Vol. 13, pp.371-382, (1982).

El-Doma, M. Age-structured Population Model with Cannibalism. Preprint to appear.

Gurtin, M. E., and R. C. MacCamy. Non-linear age-dependent Population Dynamics. *Arch. Rational Mech. Anal.* Vol. 54, pp. 281-300, (1974).

Iannelli, M., M.-Y. Kim E.-J. Park and A. Pugliese. Global boundedness of the solutions to a Gurtin-MacCamy system. *Nonlinear differ. equ. appl.* Vol. 9, pp. 197-216, (2002).

Iannelli, M. *Mathematical theory of age-structured population dynamics*. Applied mathematics monographs. Vol. 7, CNR., Giardini-Pisa, (1995).

Weinstock, E., C. Rorres. Local stability of an Age-structured Population with density-dependent fertility and mortality. *SIAM J. APPL. MATH.* Vol. 47, pp. 589-604, (1987).