



## A Complexification of Rolle's Theorem

J.P. Pemba, A.R. Davied, N.K. Muoneke

Department of Mathematics  
Prairie View A&M University  
Prairie View, TX 77446

Received July 20, 2006; revised December 14, 2006; accepted March 22, 2007

### Abstract

A new version of the classical Rolle's theorem is proved for any  $\mathbb{C}$ -valued differentiable function of the complex variable on an open connected convex subset of the complex field. The associated Mean-Value theorem follows naturally. A few explicit illustrative examples are provided in the closing section of the paper.

**Keywords:** Parametrization, simple, path-connected, smooth path

**AMS 2000 Subject Classification No.:** 30-E99, 54-F05

### 1. Introduction

Rolle's theorem together with its slant version Mean-Value theorem have represented two key ingredients in the block-by-block construction of classical Calculus of Newton and Leibnitz. Moreover, they turned out to be the main tools in the direct correlation between Analysis and Geometry so beautifully illustrated by the powerful Fundamental Theorem of Calculus. So, naturally those two theorems have been at the center of a lot of activity (Evard and Jafari (1992), ..., Tokieda Tadashi (1999)) to stretch their potential in all possible directions. Several versions have been established beyond their original real variable context into other ordered-fields (Yan Li (1999), ..., Alan Szaz (1996)) as well as some multidimensional generalizations (Jules Ferrer (1986), Furi and Martelli (1995), M.V. Korobkov (2001)). However, verbatim extensions have faced immediate problems; for example, even a simple case such as the holomorphic transcendental function  $f(z) = e^{2\pi iz} - 1$  with its infinite number of roots along the real axis, exhibits the failure of a classical Rolle's theorem in the complex variable, since its derivative cannot vanish in the complex plane.

So, more activity surrounding these two classical theorems have been centered in various types of modifications, some more elaborate than others to adapt to various existence theorems, especially in Partial Differential Equations on nonstandard domains or for many more practical reasons. In what follows, we establish some directional version of Rolle's and Mean-Value theorems in the complex field with potential implications in some Boundary-value problems with solutions sought at least in the sense of Distributions even on nonstandard domains.

## 2. Results

Evard and Jafari (1992) established the following theorem:

**Theorem 2.1. (Evard & Jafari).** Let  $f$  be a holomorphic function defined on an open convex subset  $D_f$  of  $\mathbb{C}$ . Let  $a, b \in D_f$  be such that  $a \neq b$  and  $f(a)=f(b)=0$ . Then there exists  $z_1, z_2 \in ]a, b[$  such that  $\Re(f'(z_1))=0$  and  $\Im(f'(z_2))=0$ .

So, their result requires complete analyticity of the function, a convex domain to ensure a total containment of the convex hull of every pair of points of interest, and then depart quite a bit from the classical format of Rolle's conclusion with two points where only single components of the original function satisfy local tangential flatness.

In the sequel, we return to the more conventional setting with less strenuous requirements on the function and pick a single point of directional tangency for the whole complex function with the added twist of arbitrary path connection between isopotential points.

**Theorem 2.2.** Let  $f$  represent a  $\mathbb{C}$ -valued differentiable function defined on an open path-connected subset  $\Omega$  of  $\mathbb{C}$ . Let  $a, b \in \Omega$  two distinct points such that  $f(a) = f(b)$ . Then, for any smooth simple connected path  $\Gamma \subset \Omega$  from  $a$  to  $b$  with canonical parametrization  $\gamma$ , there is some  $z_0 \in \Gamma$  such that  $f'(z_0)\gamma'(\gamma^{-1}(z_0)) \bullet (b-a) = 0$ .

**Proof:**

Let  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  be a differentiable function and  $\gamma : [0,1] \rightarrow \Gamma$  a smooth parametrization of  $\Gamma \subset \Omega$ , with  $\gamma(0) = a$  and  $\gamma(1) = b$ . Now, define a composite projection  $\Delta : [0,1] \rightarrow \mathbb{R}$  by  $\Delta(t) = f(\gamma(t)) \bullet (b-a)$ , where  $\bullet$  denotes the scalar product in the Hilbert space structure of  $\mathbb{C}$ . Clearly,  $\Delta$  satisfies all the hypotheses of the classical Rolle's theorem; Indeed,  $\Delta$  is obviously smooth on  $(0,1)$ , continuous on  $[0,1]$ , by composition and  $\Delta(0) = f(\gamma(0)) \bullet (b-a) = f(a) \bullet (b-a) = f(b) \bullet (b-a) = \Delta(1)$ . Hence, choose  $t_0 \in (0,1)$  and let  $z_0 = \gamma(t_0) \in \Gamma$  such that  $f'(z_0)\gamma'(t_0) \bullet (b-a) = 0$ .

**Remark 2.3.**

As a geometric interpretation, we can note that the image of this particular type of map is a 2D-complex manifold. Theorem 2.2 says that when two points  $a \neq b$  on that surface have the same height, then along any path (entirely contained in the surface) connecting the two points, we can find at least one point where the tangency is flat, at least in the direction of  $b - a$ .

**Theorem 2.4.** Let  $f$  represent a  $\mathbb{C}$ -valued differentiable function defined on an open path-connected subset  $\Omega$  of  $\mathbb{C}$ . Let  $a, b \in \Omega$  be two distinct points. Then, for any smooth simple connected path  $\Gamma \subset \Omega$  from  $a$  to  $b$  with canonical parametrization  $\gamma$ , there exists some  $z_0 \in \Gamma$  such that

$$\left[ f'(z_0) - \frac{f(b) - f(a)}{b-a} \right] \gamma'(\gamma^{-1}(z_0)) \bullet (b-a) = 0 .$$

**Proof:**

Given  $f : \Omega \subset \mathbb{C} \rightarrow \mathbb{C}$  satisfying the above hypotheses, we introduce

$$\Phi_f(z) = f(z) - f(a) - \frac{f(b) - f(a)}{b-a}(z-a),$$

whose  $(2 \times 2)$  Jacobian is given by

$$\begin{aligned} D\Phi_f(z) &= Df(z) - \frac{f(b) - f(a)}{b-a} \\ &\equiv \begin{bmatrix} \frac{\partial f_1}{\partial x}(z) - \Re \left\{ \frac{f(b) - f(a)}{b-a} \right\} & \frac{\partial f_2}{\partial x}(z) - \Im \left\{ \frac{f(b) - f(a)}{b-a} \right\} \\ \frac{\partial f_1}{\partial y}(z) - \Re \left\{ \frac{f(b) - f(a)}{b-a} \right\} & \frac{\partial f_2}{\partial y}(z) - \Im \left\{ \frac{f(b) - f(a)}{b-a} \right\} \end{bmatrix}, \end{aligned}$$

where  $f_1$  and  $f_2$  respectively denote the real and imaginary components of  $f$ ,  $z = x + i y$ , and such that  $\Phi_f(a) = \Phi_f(b) = 0$ . And thus, by theorem 2.2 above, we get

$$(D\Phi_f(z_0)) \gamma'(\gamma^{-1}(z_0)) \bullet (b-a) = 0 .$$

**Remark 2. 5.**

Similar to remark 2.3 above, a corresponding geometrical interpretation is made for the existence of some tangent hyperplane parallel to the support of any fixed direction through the convex hull of the manifold.

**3. Numerical Illustration**

The holomorphic function  $f(z) = e^{2\pi iz} - 1$  has zeroes at every integral value  $z = 0, \pm 1, \pm 2, \dots$ , yet its Jacobian  $f'(z) = -2\pi e^{-2\pi y} \begin{bmatrix} \sin 2\pi x & -\cos 2\pi x \\ \cos 2\pi x & \sin 2\pi x \end{bmatrix}$  never vanishes anywhere on the complex plane and thus cannot satisfy the conclusions of the classical Rolle's theorem.

On the other hand,

$\alpha)$  if  $\Gamma = [-1, 1] \times \{0\}$ , path connecting the pair of points  $a = (-1, 0)$  and  $b = (1, 0)$ , then we have  $b - a = (2, 0)$  as well as the trivial parametrization  $\gamma(t) = (2t - 1, 0)$  with  $\gamma'(t) = (2, 0) \equiv \begin{bmatrix} 2 \\ 0 \end{bmatrix}$  in matrix form. On  $\Gamma$ , we can easily solve that

$\left[ f'(x_0, 0) \gamma'(t_0) \right] \bullet \begin{bmatrix} 2 \\ 0 \end{bmatrix} = 0$  at any  $z_0 = (n, 0) \in \mathbb{Z} \times \{0\}$ ; so that the origin can be selected as solution;

$\beta$ )  $\Gamma =$  semicircle of radius 1, at the origin, as path connecting the same 2 points  $a = (-1, 0)$  and  $b = (1, 0)$ . This time, a natural parametrization could be  $\gamma(t) = (\cos \pi(1-t), \sin \pi(1-t))$  with  $\gamma'(t) = \pi(\sin \pi(1-t), -\cos \pi(1-t))$ . Then, it is easy to solve that  $z_0 = \gamma(1/2) = (0, 1)$ , and then verify that

$$\left[ f'(z_0) \gamma'(1/2) \right] \bullet \begin{bmatrix} 2 \\ 0 \end{bmatrix} = -2\pi^2 e^{-2\pi} \left\{ \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\} \bullet \begin{bmatrix} 2 \\ 0 \end{bmatrix} = 0; \text{ and}$$

$\gamma$ ) More generally, for any smooth simple connected path  $\Gamma \subset \Omega$  from  $(-1, 0)$  to  $(1, 0)$  with smooth parametrization  $\gamma$ , let  $\gamma(0) = (-1, 0)$ ,  $\gamma(1) = (1, 0)$ , and  $\gamma(t) = (x, y) = z$ ; then,  $f(\gamma(t)) = (e^{-2\pi y} \cos 2\pi x - 1, e^{-2\pi y} \sin 2\pi x)$ . We consider the special smooth real-valued map  $\Delta(t) = 2e^{-2\pi y} \cos 2\pi x$ , due to the smoothness of  $\gamma(t) = (x, y)$ . Clearly,  $\Delta(0) = \Delta(1) = 2$ ; which is enough to choose some  $t_0$  such that

$$\Delta'(t_0) = f'(\gamma(t_0)) \gamma'(t_0) \bullet \begin{bmatrix} 2 \\ 0 \end{bmatrix} = 0.$$

## REFERENCES

- Evard, J. C. and F. Jafari (1992), *A complex Rolle's theorem*, **Amer. Math. Monthly** **99** no. 9 858 – 861.
- Ferrer, J. (1986), *Rolle's theorem fails in  $l^2$* , **Amer. Math. Monthly** **93** 8 - 13.
- Furi, M. and M. Martelli (1995), *A multidimensional version of Rolle's theorem*, **Amer. Math. Monthly** **102** no. 3 243 - 249.
- Jafari, F., J. Evard, and P. Polyakov (1995), *Generalizations and applications of a complex Rolle's theorem*, **Nieuw Archief voor Wiskunde** **13** 173 - 180, **Math. Sci. Net**.
- Hansen, W. (1998), *Liouville's theorem and the restricted mean value property in the plane*, **J. Math. Pures & Appl.** **77** no. 9 943 - 947.
- Korobkov, M. V. (2001), *A generalization of the Lagrange mean value theorem to the case of vector-valued mappings*, **Siberian. Math. J.** **42** no. 2 297 - 300.
- Li, Y. (1999), *Mean value theorem for complex analytic functions*, **Natur. Sci. J. Xiangtan U.** **21** no. 4 125 - 129, **Math. Sci. Net**.
- Rosenholtz, I. (1991), *A topological mean value theorem for the plane*, **Amer. Math. Monthly** **98** 149 – 153.
- Szaz, A. (1996), *A Cauchy's mean value theorem for complex functions*, **Math. Stud.** **64** no. 1-4 125 - 127, **Math. Sci. Net**.
- Tadashi, Tokieda F. (1999), *A mean value theorem*, **Amer. Math. Monthly** **106** no. 7 673 - 674.